

# SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR OUTRIDER® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Outrider herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on this supplemental labeling and on the label pamphlet provided with the pesticide container. This supplemental labeling must be in the possession of the user at the time of pesticide application.



*Herbicide by Monsanto*

EPA Reg. No. 524-500

Outrider is a registered trademark of Monsanto Technology LLC.

## CONTROL OF JOHNSONGRASS, PURPLE AND YELLOW NUTSEdge, AND GREEN KYLLINGA IN BERMUDAGRASS AND BAHIA GRASS PASTURES

Keep out of reach of children

**CAUTION!**

In case of an emergency involving this product or for user safety information on this product, call collect, day or night (314) 694-4000.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "DIRECTIONS FOR USE" and "MIXING INSTRUCTIONS" sections of the label pamphlet for Outrider herbicide for essential product performance information.

Outrider herbicide may be used in early spring through fall to control or partially control johnsongrass, purple and yellow nutsedge, green kyllinga and other weeds in well-established bermudagrass and bahiagrass pastures. Apply Outrider herbicide up to 2.0 ounces per acre. A follow-up application can be made after suitable regrowth of weeds and no sooner than 40 days after the previous application. Use higher rates of this product for control of large, established weeds or when growth is heavy or dense. **DO NOT EXCEED TWO APPLICATIONS OR A MAXIMUM OF 2.66 OUNCES OF THIS PRODUCT PER ACRE PER YEAR.**

Addition of nonionic surfactant at 0.25 percent by volume (1 quart per 100 gallons of spray solution) is required. Use only nonionic surfactants that contain at least 90 percent active ingredient. Do not use surfactants that alter the pH of the spray solution below pH 5. Spray solutions of between pH 6.0 and 8.0 provide optimum results.

Grass forage may be grazed immediately after application. Do not harvest for hay within 14 days of application.

No crop, except wheat, may be planted into pastures that have been treated with this product within 12 months after application. All crops other than wheat may be seeded only after completion of a successful field bioassay as indicated in the FIELD BIOASSAY section.

### FIELD BIOASSAY

A field bioassay must be completed before rotating any crop other than wheat into pastures that have been treated with this product. To conduct an effective field bioassay, grow strips of the crop you intend to grow the following season in fields previously treated with Outrider herbicide. Crop response to the bioassay will determine if the crop(s) grown in test strips can be grown safely in areas previously treated with this product.

**Read the "LIMIT OF WARRANTY AND LIABILITY" in the label pamphlet for Outrider herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.**

MONSANTO COMPANY  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

©2008

4/2/2008

71014G1-19