

# SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR OUTRIDER® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Outrider herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on this supplemental labeling and on the label pamphlet provided with the pesticide container. This supplemental labeling must be in the possession of the user at the time of pesticide application.



*Herbicide by Monsanto*

EPA Reg. No. 524-500

Outrider, Roundup PRO, and Roundup PROMAX are registered trademarks of Monsanto Technology LLC.

## PASTURE AND RANGELAND SITES IN STATES WEST OF THE MISSISSIPPI RIVER

Keep out of reach of children  
**CAUTION!**

In case of an emergency involving this product or for user safety information on this product, call collect, day or night (314) 694-4000.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "DIRECTIONS FOR USE," "MIXING," and "WEEDS CONTROLLED" sections of the label pamphlet for Outrider herbicide for essential product performance information.

Outrider herbicide may be used in pasture and rangeland grasses in States west of the Mississippi River in the fall or spring to provide selective post-emergent control or partial control of the weeds specified in the "WEEDS CONTROLLED" section of the label pamphlet.

This product is selective in crested wheatgrass and selectivity in other pasture grasses is increased when they are not actively growing. Temporary stunting or chlorosis of grasses may occur but desirable grasses will recover. If concern exists about selectivity on desirable grasses, a small area should be treated to confirm selectivity.

Grass forage may be grazed immediately after application. However, for best weed control do not mow or graze the pasture or rangeland for 2 weeks before or after application.

#### Ground Broadcast and Aerial Application

Apply 0.75 to 1.33 ounces of Outrider herbicide per acre along with a nonionic surfactant. Use the higher rate when weeds are in advanced growth stage. The level of weed control following

application is dependent on weed species and weed stage of growth at application. For best results, weeds should be actively growing and in an early vegetative stage.

Refer to the "SPRAY DRIFT MANAGEMENT" section of the label pamphlet for guidelines regarding spray drift management.

#### Dormant Pastures and Rangelands

Apply 0.75 to 1.33 ounces of Outrider herbicide per acre in a tank mix with Roundup PRO® Concentrate herbicide at 10 to 13 fluid ounces per acre or Roundup PROMAX® herbicide at 9 to 11 fluid ounces per acre for control of weeds in dormant pastures. When tank mixing this product with Roundup PROMAX herbicide at rates below 12 ounces per acre, the addition of a nonionic surfactant to the spray solution at a concentration of 0.25 percent by volume (1 quart per 100 gallons of spray solution) is required. These applications should be made when the desirable pasture grass species are dormant and a new flush of the target weeds is emerged and actively growing.

**Read the "LIMIT OF WARRANTY AND LIABILITY" in the label pamphlet for Outrider herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.**

©2009 MONSANTO COMPANY  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

71014H1-20

2/2/2009